

Colonoscopy Bowel Preparation

Extra Preparation Morning (AM) Procedure

You will need to buy from your pharmacy: 4 sachets of Colonlytely (70g each) and 8 sachets of Movicol. Please disregard the instructions on the packets and follow this sheet instead.

Flushing the bowel clean of faeces is <u>very important</u> to ensure your doctor can see the inside of your colon clearly. If your colon is not clean, the procedure may have to be stopped or repeated. Please adhere to these instructions.

7 Days Before

Stop anti-diarrhoeal and iron tablets

5 Days Before

Drink 1 sachet of Movicol, morning and night (this will finish the day before your bowel preparation)

3 Days Before

Low fibre diet (see Page 2)

1 Day Before

8am: Clear fluids only

Make up each sachet of Colonlytely in 1L of water and place in the fridge.

5pm: Drink 1L Colonlytely. Continue to drink clear fluids.

6pm: Drink 1L Colonlytely. Continue to drink clear fluids.

7pm: Drink 1L Colonlytely. Continue to drink clear fluids.

Day of Colonoscopy

5am: Drink 1L Colonoscopy (this is early but very important!)

6am: You are now fasting (you may have your morning medications with a sip of water; no diabetic tablets). No food, chewing gum, lollies or drinks.

If you have any questions, please go to:

https://monashhealth.org/services/gastroenterology/patients/endoscopy/ or call the Endoscopy Bookings Office on 1800 786 188.



	Low Fibre Diet, includes:
	- white bread, white rice, pasta, noodles
10.0	- eggs (not fried), cheese
	- well-cooked vegetables (no skin)
	- chicken, fish
	- cereal
	- biscuits
	No grains, seeds, nuts, red meat, fruit, wholemeal
	bread/cereal, corn, peas, celery, carrots
	, , , , , , , , , , , , , , , , , , , ,
	Clear fluids, includes:
	- water
	- cordial
	- coffee or tea (no milk)
	- clear fruit juice (no pulp)
	- soft drink
	- clear broth
	No red or purple liquids, milk
	and the sequence, some
	Tips:
\9/	- drink clear fluids as much as possible to prevent dehydration
	(aim to drink 3-6 glasses of water whilst taking the prep)
9	- bowel prep will cause diarrhoea. Stay near a toilet.
	- finish all the prep!
L	